

**ALLERGY INJECTION PROGRAM  
PARTICIPATION CONTRACT**

**NC STATE**

2815 Cates Avenue  
Raleigh, NC 27695  
Appointments (919) 515-2563

I understand and agree to the following:

1. Allergy injections are given 8:20 am-11:00 am and 1:40 pm-3:40 pm M, W, TH, and F. (On Tuesdays they are given beginning at 9 am.) \*\*Hours may vary during school breaks and holidays. For updates, please visit <https://healthypack.dasa.ncsu.edu/>.
2. It is important to keep scheduled appointments for allergy injections. You should arrive 5-10 min. prior to your appointment time. (If you are late for your appointment, you may have to reschedule.) Please call 919-515-2563 and cancel or reschedule your appointment if you know you cannot come at your scheduled time. **\*\*Per Student Health policy- a fee is charged for all "NO SHOW" visits.**
3. You may not have more than one appointment in Student Health on the same day (ex. You cannot see a medical provider and have an allergy shot visit on the same day.)
4. Allergy injections cannot be given without complete instructions from your allergist regarding dosage, schedule of injections and treatment of possible reactions.
5. Students are responsible for ordering and receiving allergy injection vials. We will let you know when new vials need to be ordered, but **vials should be sent directly to you, NOT to Student Health.**
6. Students are responsible for retrieving and replacing their own allergy injection vials from the refrigerator in the Student Health Allergy room.
7. **Expired serum will not be used and will be discarded.**
8. Students are required to wait **30 minutes** after their allergy injection(s) are administered. Students not able to comply with this rule may be denied further treatment through the Allergy Clinic.
9. At each visit, we ask that you advise us of any change in your medical history, including pregnancy and any new medications, especially beta-blockers (used to treat blood pressure, headaches, etc.). Beta-blockers may decrease your responsiveness to epinephrine, the medicine we use to treat severe allergic reaction.
10. **If you are ill with fever, asthma or respiratory difficulties, you should not receive allergy injections until your symptoms have improved.**
11. You should not exercise or get overheated for several hours after allergy injections to minimize risk of systemic reaction.
12. You should know the signs of a systemic allergic reaction which include:

Itching of the skin or palms	Nausea/ Vomiting
Hives	Sudden Onset of hay fever symptoms
Wheezing	Flushing/ Sweating
Tightness in the throat or chest	Dizziness/ Fainting

Patient Signature \_\_\_\_\_

Date \_\_\_\_\_